

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/306239346>

Human and animal Trypanosomiasis insights from Lambwe Valley Foci, Kenya: Assessment of re-emergence potency and potential trypanotolerance

Conference Paper · August 2016

CITATIONS

0

READS

45

1 author:



[Magak A.G. Ng'wena](#)

Maseno University, School of Medicine

49 PUBLICATIONS 125 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Evaluation of Cell Surface and Serological Biomarkers for Early Diagnosis, Monitoring and Improved Management of Prostate Cancer [View project](#)



MTB/RIF [View project](#)



Human and Animal Trypanosomiasis in Lambwe Valley Foci, Kenya– Current Situation and Latent Trypanotolerance

**Ng'wena Gideon Magak¹, Willis Onyango Okoth², Owiti George Michael¹,
Onyango Kennedy¹, Owiti O. Thedeus³, Bob Awino⁴, Willis Oyieko¹,
Christopher Khayeka-Wandabwa^{5,6*} and Odero Wilson¹**

¹Maseno University School of Medicine, P.O.Box 333, Maseno, Kenya.

²Division of Vector Borne Diseases, Homa Bay County, P.O.Box 469-40300, Homa Bay, Kenya.

³Suba District Hospital, P.O.Box 25-40308, Sindo, Kenya.

⁴Maseno University, School of Public Health P.O.Box 333, Maseno, Kenya.

⁵School of Pharmaceutical Science and Technology (SPST), Health Sciences Platform,
Tianjin University, Tianjin 300072, China.

⁶African Population and Health Research Center (APHRC) P.O.Box 10787-00100, Nairobi, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. Author NGM designed the study, coordinated sample collection in the field and performed statistical analysis. Author WOO wrote the protocol for the animal studies' and performed the parasitological laboratory work. Author OGM wrote the human study protocol and coordinated the human studies. Author OK designed the human studies and wrote the protocol. Author OOT designed the human studies and coordinated the clinical assessment. BA wrote the animal studies protocol and managed literatures search. Author WO wrote the human study protocol and performed literature search. Author CKW designed the animal studies and performed statistical analysis. Author OW designed the study, coordinated sample collection and processing. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Hager Tarek Hassan Ismail, Assistant Professor, Department of Clinical Pathology, Faculty of Veterinary Medicine, Zagazig University, Egypt.

Reviewers:

(1) John Ifeanyichukwu Amaka, Kebbi State University of Science and Technology, Nigeria.

(2) Benjamin Valladares Carranza, Autonomous University of The State of Mexico, Mexico.

(3) Vincent Imieje, University of Benin, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47876>

Original Research Article

Received 18 December 2018

Accepted 09 March 2019

Published 28 March 2019

ABSTRACT

Human and animal trypanosomiasis are threats to both animal and human health in Sub-Saharan Africa. Numerous integrated vector control programs in Lambwe Valley foci have not been in tandem with insights aimed at investigating the prevalence trends besides assessing possible trypanotolerance in cattle. The cross-sectional survey enlisted human and cattle subjects for examination. Prevalence of trypanosomiasis in animal subjects was 9.2%. Blood parameters; white blood cells, lymphocytes and granulocytes were significantly lower among the parasitaemic as compared to the aparasitaemic animals except for monocyte proportions whereas, Red blood cells indices were depressed in trypanosomiasis. Viewed together, the presented findings point to animal trypanosomiasis cases in Lambwe Valley with chronic anemia and depressed white blood cell levels persisting as the most common features of the disease. High throughput molecular-based techniques currently not accessible should be incorporated into national and regional governance control programs as well as monitoring and evaluation strategies and this should be applied at both veterinary and clinical settings. Such efforts will go a long way in operationalization of modern advanced practices while mitigating likely clinical and veterinary challenge around accurate cases profiling and management based on the current widely available and applied prognosis/diagnostic strategies.

Keywords: *Trypanosomiasis; sleeping sickness; trypanotolerance; sub-saharan Africa; anaemia.*

1. INTRODUCTION

African trypanosomiasis is a vector-borne parasitic disease that affects both animal and human health in Sub-Saharan Africa. Trypanosomes, transmitted via the bite of infected *Glossina* tsetse fly as a primary vector are the responsible multi-host parasites with the capacity to infect humans, wildlife and a wide range of domestic species (that may also constitute a reservoir) [1,2]. The parasitic disease predominantly affects remote rural area communities where people get exposed to tsetse bite during their daily activities [3].

The classic human African trypanosomiasis (HAT) or sleeping sickness which is normally fatal if untreated is caused by two distinct subspecies of the African trypanosomes: *Trypanosoma brucei gambiense* (shown to account for about 95% of all reported HAT) is linked to the chronic Gambiense form of the disease in central and west Africa, with the Democratic Republic of the Congo (D.R. Congo), Central African Republic, South Sudan and Chad prevailing as the most affected countries today [2,4]. Beside the classical symptom of sleep/wake cycles reversal, the disease is also associated with anemia, immunosuppression, infertility and endocrine disorders. A person can be infected for months or even years without major signs or symptoms and when these emerge, the patient is often already in an advanced disease stage involving the central nervous system. In eastern and

southeastern Africa (including Tanzania, Malawi, Zambia and Uganda), *Trypanosoma brucei rhodesiense* is implicated in the acute Rhodesiense form of the disease with first signs and symptoms being observed a few months or weeks after infection [4,5]. Malawi and Uganda are the most affected countries with Uganda reported to support both *Trypanosoma brucei rhodesiense* and *Trypanosoma brucei gambiense* [4]. In livestock, animal African trypanosomiasis (AAT) related parasites: *Trypanosoma brucei brucei*, *Trypanosoma congolense*, *Trypanosoma vivax*, *Trypanosoma evansi* and *Trypanosoma equiperdum* have been shown to cause wasting form of the diseases resulting in major economic losses in the concerned nations [1,5]. *Trypanosoma* species that have been reported to be pathogenic to domestic as well as wild animals include *Trypanosoma congolense*, *Trypanosoma vivax* and *Trypanosoma brucei brucei* [6] whereas *Trypanosoma brucei gambiense* can infect both domestic and wild animals although their role as reservoir of the disease is contested [4]. Available reports show that sustained control efforts of the parasite and/or reduction of the vector have succeeded in lowering the number of new cases of *Trypanosoma brucei gambiense* and *rhodesiense* in quite a number of countries [5,7]. In 2009, reported human cases in Africa were about 10,000 which dropped to 7216 cases in 2012 [8-12] and in areas where the disease is endemic, *Trypanosoma brucei rhodesiense* is

apparently associated with fewer cases (less than 2% of the total reported cases).

In East Africa, trypanosomiasis has been categorized together with the neglected and possibly re-emerging zoonotic tropical diseases [13] after it was thought to be fully controlled to very low rates in the 80s. The transmitting vector, *Glossina pallidipes*, which was thought to have disappeared from the region [14], has been reported to re-invade southeastern Uganda [15] leading to significant increase in the prevalence of animals trypanosomiasis in particular. The re-invasion premise points to a contested basis for re-emergence of both human and animals' trypanosome infections due to the interconnections of the tsetse fly belts in the East African region. It has also been reported that the Lambwe vector population is more genetically diverse suggesting that these vectors have survived the numerous control efforts over the years [13]. Following the re-invasion premise and associated contentions the present study investigated the current prevalence and trend of human and animal trypanosomiasis besides assessing potential animal trypanotolerance in Lambwe valley foci.

2. METHODOLOGY

2.1 Study Design and Settings

The cross-sectional survey was conducted in the Lambwe Valley of HomaBay County, Kenya from August 2014 to April 2015. Lambwe Valley is located approximately within latitudes 0° 30' South and longitudes 34° 20' East. The area's elevation is about 1100-1400m above sea level and experiences a typical climate of two rainfall seasons with the long rains coming between March and May and short rains between September and November of every year. Daily temperatures range between 26°C to 28°C. The dominant livestock in the area includes cattle, sheep and goats and people living in the Valley undertake mixed livestock and crop farming as their main source of income. The present study targeted people and domestic animals in the catchment area of Sindo District Hospital in Lambwe Valley of Homa-Bay County within the study period. The study participants were drawn from three health facilities; Sindo district hospital, Ponge and Nyadenda dispensaries which are both located within Ruma National Game Park. The study setting is as detailed in Fig. 1. The area is one of the disease foci in East Africa and

offers a good environment for tsetse flies and wildlife which are the major reservoirs for HAT.

A sampling frame for the catchment area of focus was developed from the current (2009) National Sample Survey and Evaluation Program V (NASSEP V) frame in consultation with the local authorities. NASSRP V is a master sampling frame maintained by the Kenya National Bureau of Statistics (KNBS) and provides a basis for the implementation of household surveys in Kenya. In cognizance of the devolved system of government, NASSEP V utilizes the county boundaries and it stratifies the population into divisions, locations, sub-locations, clusters, and households. Briefly, domestic cattle in the catchment area of Sindo District Hospital and the respective feeder Ponge and Nyadenda dispensaries which are all located within Ruma National Game Park study coverage area that informed the sampling frame were sampled through consecutive sampling [16, 17]. It allowed for the inclusion of all eligible cattle subjects per eligible household within the sampling frame. Eligible households were those that keep cattle and which were identified through community leadership structure in line with the sample frame under the care of the research project enumerators duly trained. Due to the local reality of the community cultural orientation, family households exist in two subcategories: one-family households or multiple-family households. Thus in total, twenty family households keeping cattle were enrolled with between 5-30 animals per family household. For inclusion, first, eligible household head informed consent to participate in the project was requested and only those who voluntarily agreed to consent were involved in the study. Second, at the household level, all indigenous cattle of both sexes and all ages - were enrolled in the study except suckling calves and exotic breeds. Suckling calves were considered to be those aged ≤ 14 months [18, 19]. Overall, most farmers started weaning and/or stopped suckling between 6-14 months and the majority were not willing to consent suckling calves enrolment due to the perceived delicate nature of this category of cattle. Exotic breeds were excluded as they are not part of dominant cattle breeds domesticated by most farmers in the study setting per veterinary records within the local government authority. Three hundred and four (304) cattle were included in the study and assessed for signs indicative of nagana [20] by a specialized veterinary officer incorporated in the project and having experience of service in the study setting. In addition, veterinary advice was

provided to respective farmers at no costs on a measure of caution, signs to watch out for and any other nutritional and treatment guidance necessary for their livestock.

For the human subjects, fifty two (52) human participants were drawn from the aforementioned health facilities through purposive sampling [16] for the period of study. Pre-study analysis of the selected facilities past health records pointed to between 1-5 suspected cases attended to per month hence the preference for purposive sampling approach. In summary, enrolled participants were among individuals seeking treatment in the named health facilities and were recruited on the basis that they presented with signs and symptoms of trypanosomes infections [21,22]. The signs were in two categories as part of standardized clinical guidelines at the facilities;

classical symptoms: sleeping disorders (insomnia and daytime sleepiness), unexplained drowsiness, fever (that is resistant to malaria treatment), headache (that is resistant to malaria treatment), and swollen lymph nodes; other symptoms: general malaise, muscle weakness, anorexia, confusion, general body pain, chancre, restlessness, pruritus, convulsions, reduced libido (for adults), coma and stupor. Patients' examination was done by a qualified resident physician linked to the project. Patients who presented with all the classical symptoms outlined above and any one or more of the symptoms in the other symptoms category were recruited into the study following a written consent. The project did provide an option for any participant who tested positive for sleeping sickness to be put on treatment (Suramin; 20 mg/Kg body weight 5-7 days for early stage




Fig. 1. Study areas within Lambwe Valley, Homa-Bay county

cases, Melarsoprol; 3.6 mg/Kg body weight, repeated every seven days for four weeks, for late stage cases) [23] and to be followed up by the study Clinician who was based at the Sindo District Hospital.

2.2 Hematological and Parasitological Analysis

Blood samples (3-5ml) were collected from human participants into BD® vacutainers (BD Biosciences, San Jose, CA) and from animals (using sterile disposable non-pyrogenic syringes (CATHY YOUNG®, France) then transferred into heparinized vacutainer tubes) and were stored in a cool box at ambient temperature prior to being processed in the laboratory. The samples were processed for hematological indices including, hematocrit, hemoglobin (Hb), red blood cell count (RBC), white blood cell count (WBC), mean cell volume (MCV) and mean cell hemoglobin (MCH) using a Coulter counter hem analyzer (BC-3000 plus, Auto Hematology Analyzer, Mindray Co. Ltd, China). Trypanosome species detection was done by microscopy (B300, Prior Scientific Ltd) following a Giemsa staining of thin blood film preparation according to Murray et al. [24] and by buffy coat method [25].

2.3 Statistical Analysis

Reference ranges for clinical hematology were based on the guidelines in the Clinical and Laboratory Standards Institute (CLSI: formerly National Council of Clinical Laboratory Services) guideline document C28-A2 and further validated in Kenya context and in other African populations [26,27]. Conventional hematological reference ranges as earlier developed and applied were adopted for cattle [See additional file 1] [28,29]. The collected data were analyzed by a paired student's t-test or one way analysis of variance using SPSS version 20.0 software package. For all symptomatic (parasitaemic and aparasitaemic animals), emphasis was put on changes in red blood cells, hematocrit and hemoglobin for the assessment of the extent of red blood cells lysis and trypanotolerance [30-32]. Differences in mean corpuscular volume and mean corpuscular hemoglobin were used to examine general effects on the red blood cells quality while total white blood cells and the differential count values were used to assess the levels of nonspecific immunity [32,33]. For all statistical tests, $P < 0.05$ was considered significant.

3. RESULTS

The presented results point to insights of human and animal trypanosomiasis as observed in both the human and animal subjects respectively in Lambwe valley. A total of 22 male and 30 female participants were enrolled, 21% of these were aged 0-21 yrs, 29% (21-30 yrs), 25% (31-40yrs), 17% (41-50 yrs) and 8% (over 50 yrs). The majority (17) of the female participants were adults aged below 30 yrs while for the male participants 50% were aged above 30 yrs (13). All the study participants exhibited all the classical signs of trypanosomiasis while the other signs manifestation was diverse (Table 1). General malaise, muscle weakness and confusion were the most common among all the study participants. There was no study participant of either gender who was in coma or stupor during the research phase and only one case (male) with reduced libido was reported. Microscopic examination of all blood smears was negative.

Regarding the blood parameters, all the white blood cell indices were within normal range according to the clinical reference ranges (Table 2) while the RBC indices were normal except for a decline in the hematocrit in both male (35.4 ± 1.3 ; reference values 40-50%) and female (32.0 ± 0.9 ; reference values 30-50%) patients. The blood picture showed normocytic and normochromic features of the RBCs. No further analysis was carried out on the human data since the study did not detect trypanosome parasites in all the blood samples.

Of the 304 cattle that were investigated in this study, 115 (37.8%) exhibited the animal trypanosomiasis signs indicative of nagana [20] including emaciation, lacrimation and presence of swollen lymph nodes. Although the other 189 (62.2%) animals were asymptomatic, all the study animals were subjected to parasitaemia investigations. Out of the 304 animals, 28 cases were found to be parasitaemic thus suggesting a prevalence rate of 9.2%. Generally the symptomatic cattle exhibited significantly low total WBC count as compared to the asymptomatic cattle (10.1 ± 0.34 vs 11.3 ± 0.27 ; $p=0.005$) and granulocyte proportions (29.9 ± 1.16 vs 34.4 ± 0.80 ; $p=0.001$) with elevated monocyte proportion (11.2 ± 0.44 vs 9.1 ± 0.30 ; $p=0.000$) and count (1.1 ± 0.05 vs 1.0 ± 0.03 ; $p=0.016$). Furthermore, all the red blood cell indices were significantly different between the symptomatic and asymptomatic cattle except for the case of

red cell distribution width and platelets count (Table 3) while the symptomatic animals showed notable mean corpuscular volume (59.7 ± 1.91 vs 52.4 ± 0.16 ; $p = 0.000$).

When the blood indices were compared between the parasitaemic (n=28) and aparasitaemic (n=276) cattle it was observed that white blood cells (WBC) (8.7 ± 0.7 vs 11.1 ± 0.2 ; $p = 0.001$), lymphocytes (LY) (4.5 ± 0.3 vs 6.5 ± 0.2 ; $p = 0.009$) and granulocytes (GR) (2.7 ± 0.3 vs 3.7 ± 0.10 ; $p = 0.015$) were significantly lower among the parasitaemic group (Table 4). Monocyte proportions were however significantly elevated among the parasitaemic cattle as compared to the aparasitaemic group (13.3 ± 1.2 vs 9.5 ± 0.3 $p = 0.000$). All the red blood cells (RBC) indices appeared depressed except for mean corpuscular volume (MCV) (72.1 ± 7.4 vs 53.4 ± 0.2 ; $p = 0.000$) which was elevated among the parasitaemic animals. The parasitaemic animals' red blood cells hemoglobin, hematocrit, mean corpuscular hemoglobin and mean corpuscular hemoglobin concentration were significantly lower as compared to the aparasitaemic animals (Table 4). On determining the severity of anemia in the parasitaemic animals, the calculated RBC and Hb cutoff levels were $3.97 \times 10^6/\mu\text{L}$ and 7.44 g/dL while the observed levels were $3.7 \times 10^6/\mu\text{L}$ and 6.7 g/dL (Table 4), respectively which were still below the hematological references ranges applied. The presented data point to severe

anemia among all the parasitaemic animals. The blood picture showed macrocytic and normochromic RBCs. The blood platelet numbers (368.4 ± 35.1 vs 286.7 ± 6.2 ; $p = 0.000$) were significantly higher among the parasitaemic cattle as compared to the aparasitaemic group (Table 4).

4. DISCUSSION

Even though highly advanced molecular analytical approaches are highly desired [3, 4] which is a deficient operational component as the study applied standard methods available in the setting as at the time of the study, it was apparent all the study participants demonstrated classical signs for trypanosomiasis while parasitaemia assays were negative with cutting-edge molecular re-examination highly desired to validate this further [3]. Whereas 37.8% of the cattle exhibited animal trypanosomiasis signs indicative of nagana [20] including emaciation, lacrimation and presence of swollen lymph nodes while the rest were asymptomatic it would be worth noting, there is a likely underestimated transmission level due to lack of advanced routine clinical molecular based diagnostic/screening techniques at the clinical level necessary for profiling of potential cases [3-5]. In relation to *Trypanosoma* species that have been reported to be pathogenic to both domestic and wild animals [6], with their role as reservoir

Table 1. Observed symptoms and parasitaemia status in trypanosomiasis symptomatic individuals from Lambwe valley

Gender and symptoms	Male	Female
Classical signs		
Sleeping disorders (insomnia and daytime sleepiness)	22	30
Unexplained drowsiness	22	30
Fever (that is resistant to malaria treatment)	22	30
Headache (that is resistant to malaria treatment)	22	30
Swollen lymph nodes	22	30
Other signs		
General malaise	15	20
Muscle weakness	10	13
Anorexia	5	8
Confusion	0	3
General body pain	9	12
Chancre	0	3
Restlessness	1	4
Pruritus	0	2
Convulsions	0	5
Reduced libido (for adults)	1	0
Coma	0	0
Stupor	0	0
Parasitaemia	Nil	Nil

Table 2. Observed Means±SEM of white blood cell, red blood cell and platelet indices of trypanosomiasis symptomatic individuals from Lambwe Valley

Gender and Indices	Males (N=22)	Female (N=30)
a) White blood cell indices		
White blood cells (WBC) $\times 10^3/\mu\text{L}$	7.8±0.8	6.9±0.5
Lymphocytes (LY) [%]	38±3.6	37±3.0
Monocyte (MO) [%]	8.3±0.9	7.6±0.9
Granulocytes (GR) [%]	53.1±3.8	53.5±3.3
Lymphocytes (LY) $\times 10^3/\mu\text{L}$	3.3±0.6	2.8±0.4
Monocyte (MO) $\times 10^3/\mu\text{L}$	0.6±0.09	0.5±0.06
Granulocytes (GR) $\times 10^3/\mu\text{L}$	3.8±0.5	3.7±0.3
b) Red blood cells indices		
Red blood cells (RBC) $\times 10^6/\mu\text{L}$	4.7±0.2	4.3±0.2
Hemoglobin (Hb) [g/dL]	12.3±0.4	11.0±0.3
Hematocrit (Hct) [%]	35.4±1.3	32.0±0.9
Mean Corpuscular Volume (MCV) [fL]	74.9±3.2	76.4±2.7
Mean Corpuscular Hemoglobin (MCH) [pg]	26.0±1.0	26.2±0.9
Mean corpuscular Hemoglobin Concentration (MCHC) [g/dL]	34.8±0.3	34.4±0.2
Red cell distribution width (RDW) [%]	15.7±0.8	16.0±0.9
Platelet (PLT) $\times 10^3/\mu\text{L}$	202.4±19.2	207.0±17.4
Mean platelet volume (MPV) [fL]	7.3±0.3	7.6±0.2

Table 3. Observed Means±SEM of white blood cell, red blood cells and platelets indices of trypanosomiasis symptomatic and asymptomatic cattle

Health status and Indices	Asymptomatic cattle (N=189)	Symptomatic cattle (N=115)	p-value
a) White blood cell indices			
White blood cells (WBC) $\times 10^3/\mu\text{L}$	11.3±0.27	10.1±0.34	0.005
Lymphocytes (LY) [%]	56.6±0.80	57.7±1.03	0.388
Monocyte (MO) [%]	9.1±0.30	11.2±0.44	0.000
Granulocytes (GR) [%]	34.4±0.80	29.9±1.16	0.001
Lymphocytes (LY) $\times 10^3/\mu\text{L}$	6.7±0.32	5.8±0.24	0.068
Monocyte (MO) $\times 10^3/\mu\text{L}$	1.0±0.03	1.1±0.05	0.016
Granulocytes (GR) $\times 10^3/\mu\text{L}$	3.8±0.1	3.3±0.28	0.084
b) Red blood cells indices			
Red blood cells (RBC) $\times 10^6/\mu\text{L}$	5.7±0.04	4.3±0.06	0.000
Hemoglobin (Hb) [g/dL]	10.6±0.09	8.4±0.13	0.000
Hematocrit (Hct) [%]	29.9±0.19	24.8±0.12	0.000
Mean Corpuscular Volume (MCV) [fL]	52.4±0.16	59.7±1.91	0.000
Mean Corpuscular Hemoglobin (MCH) [pg]	18.4±0.13	19.6±0.21	0.000
Mean corpuscular Hemoglobin Concentration (MCHC) [g/dL]	35.4±0.23	33.9±0.50	0.003
Red cell distribution width (RDW) [%]	35.3±0.53	36.2±0.55	0.238
Platelets (PLT) $\times 10^3/\mu\text{L}$	289.4±7.60	302.1±12.14	0.352

being contested [4], it would be of interest for further research in sub-Saharan context to aim at monitoring the African trypanosomiasis epidemiologic patterns in a consistent trend. The monitoring efforts should target complete eradication without underestimating low transmission burden for every life towards better health matters in tune with the sustainable development goals three[3, 34].

The main intervention approaches against sleeping sickness have been active and passive case-finding for early detection of cases followed by treatment, vector control and animal reservoir management. This should be continued alongside proper disease surveillance programs anchored on modernized molecular diagnostic approaches at the most localized primary health

Table 4. White blood cells, red blood cells and platelets indices of trypanosomiasis parasitaemic and aparasitaemic cattle cases

Trypanosomiasis status and Indices	-VE (N=276)	+VE (N=28)	p-value
a) White blood cell indices			
White blood cells (WBC) $\times 10^3/\mu\text{L}$	11.1 \pm 0.2	8.7 \pm 0.7	0.001
Lymphocytes (LY) [%]	57.3 \pm 0.6	53.7 \pm 2.5	0.091
Monocyte (MO) [%]	9.5 \pm 0.3	13.3 \pm 1.2	0.000
Granulocytes (GR) [%]	32.9 \pm 0.7	30.7 \pm 2.3	0.351
Lymphocytes (LY) $\times 10^3/\mu\text{L}$	6.5 \pm 0.2	4.5 \pm 0.3	0.009
Monocytes (MO) $\times 10^3/\mu\text{L}$	1.0 \pm 0.00	1.1 \pm 0.1	0.359
Granulocytes (GR) $\times 10^3/\mu\text{L}$	3.7 \pm 0.10	2.7 \pm 0.3	0.015
b) Red blood cells indices			
Red blood cells (RBC) $\times 10^9/\mu\text{L}$	5.3 \pm 0.04	3.7 \pm 0.2	0.000
Hemoglobin (Hb) [g/dL]	10.1 \pm 0.08	6.7 \pm 0.3	0.000
Hematocrit (Hct) [%]	28.3 \pm 0.2	24.2 \pm 0.3	0.000
Mean Corpuscular Volume (MCV) [fL]	53.4 \pm 0.2	72.1 \pm 7.4	0.000
Mean Corpuscular Hemoglobin (MCH) [pg]	18.9 \pm 0.1	18.1 \pm 0.3	0.037
Mean corpuscular Hemoglobin Concentration (MCHC) [g/dL]	35.6 \pm 0.2	27.5 \pm 1.0	0.000
Red cell distribution width (RDW) [%]	35.8 \pm 0.4	34.2 \pm 0.8	0.234
Platelets (PLT) $\times 10^3/\mu\text{L}$	286.7 \pm 6.2	368.4 \pm 35.1	0.000

care level (health centers in Kenya) that serve the majority of the affected locals. Therefore, successful strategies for controlling African trypanosomiasis should be pegged onto the precision of techniques applied at early diagnosis and treatment, as well as vector control. Land use and land cover should be adopted for vector eradication as the communities in Lambwe Valley also modify their cattle grazing and watering points in order to reduce the animals' interaction with the vectors. In addition, National and Regional Control Initiatives such as by the Pan African Tsetse and Trypanosomiasis Eradication Program (PATTEC) should be greatly enhanced to continue improving human and animal health in most parts of Lambwe Valley through integrated vector control and follow-ups.

Trypanosomiasis pathology is mainly associated with the disease's effects on blood and particularly the fact that the trypanosomes parasitize the RBCs in addition to invading the CNS [35, 36]. The depressed levels of WBC, LY and GR in all the parasitaemic cattle in this study were suggestive of chronic parasitic infection. Due to the trypanosome parasites' antigenic variations abilities and host immune systems evasion, the disease is usually associated with bouts of changes in white blood cell numbers and proportions [6, 37]. The WBC numbers will tend to increase [38] at the onset but decrease with time depending on the duration of the infection. The leucopenia and especially

lymphopenia in the study animals may have been compounded by the infection status and anorexia nervosa as the animals appeared emaciated and lethargic. All the parasitaemic animals would also have suffered the deficiency of some of the hemopoietic requirements such as depressed endocrine, nutrients and cytokine factors [32].

The severe anemia in cattle reported in this study is suggestive of the fact that the animals in Lambwe valley may not be trypanotolerant. From previous reports severity of anemia has been used to measure the domestic animal's tolerance to trypanosomes infection. This is because anemia is recognized as the main consequence of trypanosomiasis and is a reliable indicator of the disease [39]. It has also been reported that the severity of anemia correlates well with the productive performance of trypanosome-infected cattle [40, 41]. Previous studies have reported that some breeds of cattle, sheep and goats have the capacity to develop less severe anemia during trypanosomes' infection. Such animals are regarded as trypanotolerant and may be used to improve animal productivity in areas such as Lambwe Valley. The MCV was however higher in the parasitaemic animals and the blood picture was that of microcytic and normochromic RBCs. These findings show that the parasitaemic animals exhibited normochromic anemia associated with leucopenia [42] which attributed such outcomes to the cleavage of erythrocyte

sialic acid of cattle infected with *T. congolense*. The RBCs in such a case become prone to phagocytosis by mononuclear phagocyte system in addition to parasitization from the trypanosomes.

The study findings presented in the context of real-world clinical protocols at the study site remain alive to the sensitivity debate of the techniques applied as a limitation particularly in human cases profiling and absence of anaplasma assessment. Incorporating additional molecular screening approaches was a constraint due to limited resources to mobilize and operationalize the same. Despite robust molecular diagnostics having been developed in the recent past, few are yet to reach the patients at primary point of care or attain incorporation into the national control programs whereas the disease burden is prominent in mostly remote settings. Of note, this limitation among other realities is not just a challenge for the current context and study but a discourse of intense debate between the neglected tropical disease support and financing agencies vis-a-vis science and public health experts as observed [3-5, 7].

5. CONCLUSION

The findings point to animal trypanosomiasis cases in the Lambwe Valley with chronic anemia and low white blood cell levels persisting as the most common clinical features. There is a need to mitigate likely underestimation of human and veterinary trypanosomiasis case burden based on the current, widely available and applied clinical screening/diagnostic strategies. Therefore, high throughput molecular based screening/diagnostic techniques currently not accessible should be incorporated into national and regional governance control programs as well as monitoring and evaluation strategies and be applied at both veterinary and human clinical medicine settings. It would be worthwhile for intervention partners' efforts in Lambwe valley to attempt focusing towards pilot research programs on the viability of trypanotolerant domestic animals for sustainable one health.

CONSENT AND ETHICAL APPROVAL

This study received Ethical approval from the Institutional Review Ethics Committee (IREC), Moi Teaching and Referral Hospital (MTRH) Kenya. Written informed consent for voluntary participation was obtained from all human participants. Further permission to get blood

samples from domestic animals was obtained from the owners.

ACKNOWLEDGEMENT

We thank Mr. George Otieno Ondata, the laboratory technologist who prepared and read all the blood slides in addition to carrying out the full hemogram assessments. This paper was published with permission from the Dean School of Medicine, Maseno University. This study was supported by funds from Partnerships for innovative Medical Education Kenya (PRIME-K).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anderson NE, et al. Characterisation of the wildlife reservoir community for human and animal trypanosomiasis in the Luangwa Valley, Zambia. *PLoS Neglected Tropical Diseases*, 2011;5(6):e1211.
2. Truc P, et al. Atypical human infections by animal trypanosomes. *PLoS Neglected Tropical Diseases*. 2013;7(9):e2256.
3. Molyneux D, Ndung'u J, Maudlin I. Controlling sleeping sickness—"when will they ever learn?". *PLoS Neglected Tropical Diseases*. 2010;4(5):e609.
4. Büscher P, Deborggraeve S. How can molecular diagnostics contribute to the elimination of human African trypanosomiasis? *Expert Review of Molecular Diagnostics*. 2015;15(5):607-615.
5. Aksoy S, et al. Human African trypanosomiasis control: Achievements and challenges. *PLoS Neglected Tropical Diseases*. 2017;11(4):e0005454.
6. Abenga JN, Anosa VO. Clinical studies on experimental Gambian trypanosomiasis in vervet monkeys. *Veterinarski Arhiv*. 2006;76(1):11-18.
7. Meyer A, et al. Past and ongoing tsetse and animal Trypanosomiasis control operations in five African countries: A systematic review. *PLoS Neglected Tropical Diseases*. 2016;10(12):e0005247.
8. Simarro PP, et al. The human African trypanosomiasis control and surveillance programme of the World Health

- Organization 2000–2009: the way forward. *PLoS Negl Trop Dis.* 2011;5(2):e1007.
9. Chappuis F, et al. Human African trypanosomiasis in areas without surveillance. *Emerging Infectious Diseases.* 2010;16(2):354-357.
 10. Cattand P, Jannin J, Lucas P. Sleeping sickness surveillance: An essential step towards elimination. *Tropical Medicine & International Health.* 2001;6(5):348-361.
 11. Organization WH. Human African trypanosomiasis (sleeping sickness): epidemiological update. *Wkly Epidemiol Rec.* 2006;81(8):71-80.
 12. Franco JR, et al. *Epidemiology of human African trypanosomiasis*; 2015.
 13. Ouma JO, et al. Genetic diversity and population structure of *Glossina pallidipes* in Uganda and western Kenya. *Parasit Vectors.* 2011;4:122.
 14. Waiswa C, et al. *Glossina fuscipes fuscipes* in the trypanosomiasis endemic areas of south eastern Uganda: Apparent density, trypanosome infection rates and host feeding preferences. *Acta Tropica.* 2006;99(1):23-29.
 15. Magona J, et al. Implications of the re-invasion of Southeast Uganda by *Glossina pallidipes* on the epidemiology of bovine trypanosomiasis. *Veterinary Parasitology.* 2005;128(1):1-9.
 16. Martínez-Mesa J, et al. Sampling: How to select participants in my research study? *Anais Brasileiros de Dermatologia.* 2016;91(3):326-330.
 17. Omair A. Sample size estimation and sampling techniques for selecting a representative sample. *Journal of Health Specialties.* 2014;2(4):142.
 18. Enríquez D, Hötzel MJ, Ungerfeld R. Minimising the stress of weaning of beef calves: A review. *Acta Veterinaria Scandinavica.* 2011;53(1):28.
 19. Das S, Redbo I, Wiktorsson H. Effect of age of calf on suckling behaviour and other behavioural activities of Zebu and crossbred calves during restricted suckling periods. *Applied Animal Behaviour Science.* 2000;67(1):47-57.
 20. Eshetu E, Begejo B. The current situation and diagnostic approach of Nagana in Africa: A review. *Journal of Natural Sciences Research.* 2015;5(17):117-125.
 21. Palmer JJ, et al. Syndromic algorithms for detection of gambiense human African trypanosomiasis in South Sudan. *PLoS Neglected Tropical Diseases.* 2013;7(1):e2003.
 22. Brun R, et al. Human african trypanosomiasis. *The Lancet.* 2010; 375(9709):148-159.
 23. Kagira J, et al. Prevalence and types of coinfections in sleeping sickness patients in Kenya (2000/2009). *Journal of tropical medicine,* 2011. 2011.
 24. Murray M, Murray P, McIntyre W. An improved parasitological technique for the diagnosis of African trypanosomiasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene.* 1977;71(4): 325-326.
 25. Murray M. *Livestock productivity and trypanotolerance: network training manual.* 1983: ILRI (aka ILCA and ILRAD).
 26. Kibaya RS, et al. Reference ranges for the clinical laboratory derived from a rural population in Kericho, Kenya. *PLoS one.* 2008;3(10):e3327.
 27. Karita E, et al. CLSI-derived hematology and biochemistry reference intervals for healthy adults in eastern and southern Africa. *PLoS one.* 2009;4(2):e4401.
 28. Latimer KS, Duncan. *Prasse's veterinary laboratory medicine: Clinical pathology.* 2011: John Wiley & Sons.
 29. Thrall MA, et al. *Veterinary hematology and clinical chemistry.* 2012: John Wiley & Sons.
 30. Murray M. Trypanotolerance, its criteria and genetic and environmental influences. *Livestock Production in Tsetse Affected Areas of Africa,* Eds ILCA/ILRAD. 1988; 133-151.
 31. Murray M, Trail J, d'Ieteren G. Trypanotolerance in cattle and prospects for the control of trypanosomiasis by selective breeding. *Rev Sci Tech.* 1990;9(2):369-386.
 32. Naessens J. Bovine trypanotolerance: A natural ability to prevent severe anaemia and haemophagocytic syndrome? *International Journal for Parasitology.* 2006;36(5):521-528.
 33. Verhulst A, Pandey V. Technical problems in the estimation of trypanotolerance within cattle and small ruminant breeds. *Proceedings of the FAO Expert Consultation on the Genetic Aspects of Trypanotolerance,* Rome, Italy, 3-4 September 1991, 1992.

34. Nam UV. Transforming our world: The 2030 agenda for sustainable development; 2015.
35. Mutayoba B, et al. Comparative trypanotolerance of the Small East African breed of goats from different localities to *Trypanosoma congolense* infection. *Veterinary Parasitology*. 1989;31(2):95-105.
36. Pentreath VW, et al. Pathogenesis of human African trypanosomiasis. *The Trypanosomiasis*. 2004;283-301.
37. Ochiogu I, et al. Experimental *Trypanosoma brucei* infection in rats (*Rattus norvegicus*): effects on different stages of gestation and the neonatal period. *Animal Reproduction*. 2008;5:39-49.
38. Turay A, et al. A comparative study on the susceptibility of male and female albino mice to *Trypanosoma brucei brucei*. *Journal of Vector Borne Diseases*. 2005;42(1):15.
39. Murray P, et al. Trypanosomiasis in N'Dama and Zebu cattle. The influence of weight of infection on the severity of the disease. 1979.
40. Henson J. The International Laboratory for Research on Animal Diseases [Kenya, East Africa, trypanosomiasis, theileriosis]. in Workshop on theileriosis. Nairobi (Kenya). 7 Dec 1976. 1977.
41. Ackah P, et al. The ILCA/ILRAD Trypanotolerance Network: situation report, December 1985, proceedings of a network meeting held at ILCA, Nairobi; 1986.
42. Esievo K, et al. Variation in erythrocyte surface and free serum sialic acid concentrations during experimental *Trypanosoma vivax* infection in cattle. *Research in Veterinary Science*. 1982;32(1):1-5.

Additional file 1. Hematologic reference ranges

	Conventional (USA) Units	SI Units	Dog	Cat	Cow	Horse	Pig	Sheep	Goat	Rabbit	Llama	Vietnamese Potbellied Pig	Ostrich
	Conventional (USA) Units	SI Units	Dog	Cat	Cow	Horse	Pig	Sheep	Goat	Rabbit	Llama	Vietnamese Potbellied Pig	Ostrich
PCV	%	$\times 10^{-2}$ L/L	35–57	30–45	24–46	27–43	36–43	27–45	22–38	33–50	29–39	22–50	32
Hgb	g/dL	$\times 10$ g/L	11.9–18.9	9.8–15.4	8–15	10.1–16.1	10–16	9–15	8–12	10–17	12.8–17.7	7.8–16.2	12.2
RBCs	$\times 10^6/\mu\text{L}$	$\times 10^{12}/\text{L}$	4.95–7.87	5.0–10.0	5.0–10.0	6.0–10.4	5–8	9–15	8–18	5–8	11.3–17.7	3.6–7.8	1.7
Reticulocytes	%	%	0–1.0	0–0.6			0–0.1						
Absolute reticulocyte count	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	<80	<60									
MCV	fL	fL	66–77	39–55	40–60	37–49	50–68	28–40	16–25	58–67	20.9–28	55–71	174
MCH	pg	pg	21.0–26.2	13–17	11–17	13.7–18.2	17–21	8–12	5.2–8			18–24	61
MCHC	g/dL	$\times 10$ g/L	32.0–36.3	30–36	30–36	35.3–39.3	30–34	31–34	30–36	29–37	43.1–46.6	31–36	33
Platelets	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	211–621	300–800	100–800	117–256	200–500	800–1,100	300–600	250–650		204–518	
MPV	fL	fL	6.1–10.1	12–18	12–18	3.5–6.5	4.0–6.0						
WBCs	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	5.0–14.1	5.5–19.5	4.0–12.0	5.6–12.1	11–22	4–8	4–13	5.2–12.5	7.5–21.5	5.2–17.9	5.5
Neutrophils (segmented)	%	%	58–85	45–64	15–33	52–70	28–47	10–50	30–48	20–75	60–74	0–63	63 ^a
Neutrophils (band)	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	2.9–12.0	2.5–12.5	0.6–4.0	2.9–8.5	2–15	0.7–6.0	1.0–7.2	1–9.4	4.6–16	0–11.4	3.4
Lymphocytes	%	%	0–3	0–2	0–2	0–1	0–4	0	rare		0–1	0–1	
Monocytes	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	0–0.45	0–0.3	0–0.1	0–0.1	0–0.8				0–0.35	0–0.19	
Eosinophils	%	%	8–21	27–36	45–75	21–42	39–62	40–55	50–70	30–85	13–35	15–55	34
Basophils	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	0.4–2.9	1.5–7.0	2.5–7.5	1.2–5.1	3.8–16.5	2–9	2–9	1.6–10.6	1–7.5	0.8–9.8	1.87
M:E	%	%	2–10	0–5	0–8	0–6	2–10	0–6	0–4	1–4	1–4	0–13	2.8
Plasma proteins	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	0.1–1.4	0–0.9	0–0.9	0–0.7	0–1	0–0.75	0–0.55	0.05–0.5	0.05–0.8	0–0.67	0.15
Plasma fibrinogen	%	%	0–9	0–4	0–20	0–7	0.5–11	0–10	1–8	1–4	0–15	0–12	0.3
	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	0–1.3	0–0.8	0–2.4	0–0.8	0–1.5	0–1	0.05–0.65	0.05–0.5	0–3.3	0–0.73	0.02
	%	%	0–1	0–1	0–2	0–2	0–2	0–3	0–1	1–7	0–2	0	0.2
	$\times 10^3/\mu\text{L}$	$\times 10^9/\text{L}$	0–0.14	0–0.2	0–0.2	0–0.3	0–0.5	0–0.3	0–0.12	0.05–0.9	0–0.4	0–0.61	0.01
	g/dL	10 g/L	0.75–2.5	0.6–3.9	0.3–1.8	0.5–1.5	1.2–2.2	0.77–1.7	0.7–1.0			5.4–7.5	
	mg/dL	g/L	6.0–7.5	6.0–7.5	6.0–8.0	6.0–8.5	6–8	6–7.5	6–7.5	5.4–8.3	200–400	100–400	100–400

Data on various species compiled and adapted in part from multiple sources, including Latimer KS, Duncan & Prasse's *Veterinary Laboratory Medicine: Clinical Pathology*, 5th ed., Wiley-Blackwell, 2011; and Weiss DJ, Wardrop KJ, *Schalm's Veterinary Hematology*, 6th Ed., Wiley-Blackwell, 2010. Reference ranges vary between laboratories. Values provided by the reference laboratory should be used.

MCV = mean corpuscular volume, MCH = mean corpuscular hemoglobin, MCHC = mean corpuscular hemoglobin concentration, MPV = mean platelet volume, M:E = myeloid:erythroid ratio^a Heterophil

© 2019 Magak et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47876>